

TWINN CROP TRIAL



Irrigated Rice, Moshi, Tanzania, 2008

KEY RESULT

A single application of TwinN plus 50% of normal N application (100 kg urea) showed no significant difference in yield compared to the standard N treatment (200 kg urea) in an independent replicated trial. TwinN applied without any added N produced a decreased yield compared to the standard N treatment.

TREATMENTS

- T1 Control (no fertilizer material applied)
T2 FYM (Farm Yard Manure) alone (10t/ha)
T3 FYM (10t/ha) + 50% recommended N from urea (100 kg/ha)
T4 TwinN alone (recommended rate)
T5 TwinN (recommended rate) + 50% recommended N from urea (100 kg/ha)
T6 Azolla (500 kg/ha) alone
T7 Azolla (500 kg/ha) + 50% recommended N from urea (100 kg/ha)
T8 100% recommended N from urea (200 kg/ha)
- Rice Cultivar: IR 64

TRIAL RESULTS: Grain Yield & Mean Values of Other Parameters

Treatment	Yield T/ha	Panicles/m ²	Grain No. per Panicle	1000 Grain Wt (g)	Plant Height (cm)	No. of Tiller/hill	Ripening Ratio (%)
T1	3.8	329.2	64.7	64.7	74.6	15.3	71.6
T2	4.0	329.2	65.8	65.8	75.0	16.0	70.3
T3	5.5	350.8	81.8	81.8	77.7	17.3	63.6
T4	4.0	330.0	62.4	62.4	75.5	13.3	70.6
T5	6.0	400.8	84.4	84.4	81.2	21.7	59.3
T6	3.8	323.3	68.9	68.9	76.8	16.0	63.4
T7	5.2	413.3	67.5	67.5	79.0	17.7	63.9
T8	6.2	455.0	87.6	87.6	83.7	22.0	55.8
CV(%)	16.2	12.1	10.5	9.8	4.5	17.8	8.7
F-Test	*	*	**	ns	ns	*	*
LSD 5%	1.0	63.8	11.0	-	-	4	8.1
LSD 1%	1.5	95.1	16.0	-	-	7	4

* Significance at 5% in F-test **Significance at 1% in F-test ns = not significant.

Mapleton Agri Biotech Pty Ltd

137 Obi Obi Road, Mapleton Qld 4560 Australia

Phone: +61 7 5445 7151 Fax: +61 7 5445 7769
Email: info@mabiotech.com www.mabiotech.com

Mapleton International Pty Ltd

Phone: +44 1666 849415
Email: info@mapletoninternational.com

www.mapletoninternational.com

Your Local Distributor

The highest yield (6.2 t/ha) was obtained in plots which received a recommend rate of N fertiliser (T-8, 200 kg urea/ha) from a chemical fertiliser (T-8). This was not significantly different to plots in which TwinN was sprayed and supplemented by 50% recommended rate of nitrogen from urea (T-5, 6.0 t/ha). FYM application at 10t/ha with fertiliser at 100kg N/ha (T-3) came third in terms of yield (5.5 t/ha). TwinN alone resulted in decreased yield compared to the full rate of N application.

TRIAL SUMMARY

Trial Performed & Analysed By: Crop Science and Production Section, KATC-Moshi
Trial Design: RCBD design, with 3 replications, 8 treatments

CONCLUSIONS

- ♦ Application of TwinN to rice enabled a 50% reduction in urea application to produce comparable yield to 200 kg urea/ha (100% rate).
- ♦ Application of TwinN with no urea did not show a significant yield increase indicating that it should be applied with either some nitrogen fertiliser or an alternative source of organic nitrogen.



Mapleton Agri Biotec Pty Ltd

137 Obi Obi Road, Mapleton Qld 4560 Australia

Phone: +61 7 5445 7151 Fax: +61 7 5445 7769
Email: info@mabiotec.com www.mabiotec.com

Mapleton International Pty Ltd

Phone: +44 1666 849415
Email: info@mapletoninternational.com

www.mapletoninternational.com

Your Local Distributor